# IBUPROFEN- ibuprofen tablet, film coated Major Pharmaceuticals

-----

## Major Pharmaceuticals Ibuprofen Tablets, 200mg Drug Facts

## Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

## **Purposes**

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

- or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

## Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

## When using this product

• take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used

## Adults and children 12 years and older:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

## Children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

## **Inactive ingredients**

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

#### Questions or comments?

1-800-616-2471

#### **Principal Display Panel**

SEE NEW WARNINGS

Ibuprofen Tablets, 200mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

COMPARE TO the active ingredient of ADVIL®

50 COATED CAPLETS\*\*

\*\* Capsule Shaped Tablets

Actual Size



KEEP CARTON FOR REFERENCE

DONOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

#### Drug Facts

Purposes Active ingredient (in each caplet) lbuprofen 200 mg (NSAID)\*...... ....Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis

■ toothache ■ backache ■ the common cold

menstrual cramps temporarily reduces fever

#### Warnings

Allergy allert: Ibup rofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs,

stop use and seek medical help right away. Stomach bleeding warming: This product contains an NSAID, which may cause severe stornach bleeding. The chances are higher if you ■ are age 60 or older ■ have had stomach ubers or bleeding problems take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

He art attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The riskis higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to ibup rofen or any other pain reliever/fever reducer in right before or after heart surgery

#### Drug Facts (continued)

Ask a doctor before use if ■ you have problems or serious side effects from taking pain refevers or fever reducers 🔳 the stomach bleeding warning applies to you who you have a history of stomach problems, such as heartburn ■ yo ú have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin under a doctor's care for any serious condition ■taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and as ka doctor if wovou experience any of the following signs of stomach bleeding: we feel faint wovemit blood whave bloody or black stools whave stomach pain that does not get better ■ you have symptoms of he art problems or stroke: ■ chest pain

mtrouble breathing mweakness in one part or side of body muslurred speech mileg swelling main gets worse or lasts more than 10 days ■fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed todo so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right a way (1-800-222-1222).

Directions • do not take more than directed ■ the small est effective dose should be used

Distrib uted by: MAJOR® PHARMACEJTICALS 17177 NLaurel Park Drive, Suite 233 Livonia, MI 48152 REV. 09'17 Re-Order No. 700128

Inactive Ingredients coloidalsilion nidoide, com starch, croscarmellose sodium, hypromellose, iron oidered, iron oide jellow, microsystelline cellulose, polyethyte neiglycd, polysob ate 80 Auestions or comments? 1-800-616-2471 teaic acid, titan ium dioxide

Adults and children 12 years and older: take
4 to 6 hours while symptoms peasist in pain or it
respond to 1 caplet, 2 cap letsmay be used ind on
6 caplets in 24 hours, unless directed by a doctor
Children under 12 years: aska dod or Other Information  $\blacksquare$  read all wan higher directions between  $\blacksquare$  store at 20-25  $\heartsuit$  (68-77°F)  $\blacksquare$  avoid high humility and excessive heat above 40  $\heartsuit$  (104°F) Ipain orfener does no ed ∎d om otexceed e ke 1 cap let ever

Drug Facts (continued)

## **IBUPROFEN**

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7912
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPRO FEN (UNII: WK2XYI10 QM) (IBUPRO FEN - UNII: WK2XYI10 QM)	IBUPROFEN	200 mg	

Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)		
STARCH, CORN (UNII: O8232NY3SJ)		
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POLYSORBATE 80 (UNII: 6 OZP39 ZG8 H)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		

Product Characteristics			
Color	BROWN	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	12
Contains			

P	Packaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0904-7912-51	1 in 1 CARTON	08/16/1994	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0904-7912-59	1 in 1 CARTON	08/30/1994	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0904-7912-24	1 in 1 CARTON	06/17/1998	09/07/2015
3		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA072096	08/16/1994

## Labeler - Major Pharmaceuticals (191427277)

Revised: 8/2020 Major Pharmaceuticals